

The
"Entwurf"
Paper

S1 Equations of motion of the material point in the static gravitational field

Free fall in a static field

$$0 = \delta \left(ds = \delta \left(-dx^2 - dy^2 - dz^2 + c^2 (1, y, z) dt^2 \right) \right)$$

$$= \delta \left(\underbrace{\sqrt{c^2 - \dot{x}_i \dot{x}_i}}_{\text{"H"}} dt \right)$$

$x_i = x, y, z$
summation of repeated i

where 1912 theory concluded

Euler-Lagrange equation

$$\frac{\partial H}{\partial x_i} - \frac{d}{dt} \frac{\partial H}{\partial \dot{x}_i} = 0 \text{ with:}$$

$$\frac{\partial H}{\partial x_i} = \frac{\partial}{\partial x_i} (c^2 - \dot{x}_i \dot{x}_i)^{\frac{1}{2}} = \frac{1}{2} \frac{1}{\sqrt{c^2 - \dot{x}_i \dot{x}_i}} \cdot \frac{\partial c^2}{\partial x_i} = \frac{c \frac{\partial c}{\partial x_i}}{\sqrt{c^2 - \dot{x}_i \dot{x}_i}}$$

$$\frac{\partial H}{\partial \dot{x}_i} = \frac{\partial}{\partial \dot{x}_i} (c^2 - \dot{x}_i \dot{x}_i)^{\frac{1}{2}} = \frac{1}{2} \frac{1}{\sqrt{c^2 - \dot{x}_i \dot{x}_i}} \cdot \frac{\partial \dot{x}_k \dot{x}_k}{\partial \dot{x}_i} = \frac{\dot{x}_i}{\sqrt{c^2 - \dot{x}_k \dot{x}_k}}$$

conserved Hamiltonian

$$E = J_i \dot{x}_i - mH$$

add m

$$\frac{d}{dt} \left\{ \frac{m \dot{x}_i}{\sqrt{c^2 - \dot{x}_k \dot{x}_k}} \right\} = - \frac{m c \frac{\partial c}{\partial x_i}}{\sqrt{c^2 - \dot{x}_k \dot{x}_k}}$$

Gravitational force

canonical momentum $J_i = m \frac{\partial H}{\partial \dot{x}_i} = \frac{m \dot{x}_i}{\sqrt{c^2 - \dot{x}_k \dot{x}_k}} \approx \frac{m \dot{x}_i}{c}$
small velocities

$$E = \frac{m \dot{x}_i \dot{x}_i}{\sqrt{c^2 - \dot{x}_k \dot{x}_k}} - \frac{m(c^2 - \dot{x}_i \dot{x}_i)}{\sqrt{c^2 - \dot{x}_k \dot{x}_k}} = \frac{m c^2}{\sqrt{c^2 - \dot{x}_k \dot{x}_k}} \approx m c + \frac{1}{2} m \dot{x}_k \dot{x}_k$$

same as $m \sqrt{c^2 - \dot{x}_k \dot{x}_k}$ small velocities

§2 Equations of motion ... arbitrary gravitational field

Particle in free fall follows $\int ds = 0$ where

$$ds^2 = \sum_{\mu\nu} g_{\mu\nu} dx^\mu dx^\nu$$

↑
"fundamental tensor"

standard analysis in Lagrangian mechanics (as before)

Canonical momentum $J_x = -\sum_\nu m g_{1\nu} \frac{dx^\nu}{ds}$ x_1, x_2, x_3, x_4
 $\nu = 1, 2, 3, 4$

Force $K_x = -\frac{1}{2} m \sum_{\mu\nu} \frac{\partial g_{\mu\nu}}{\partial x_1} \frac{dx^\mu}{ds} \cdot \frac{dx^\nu}{dt} \Big|_{x_4}$

Special case of static field

$$g_{\mu\nu} = \begin{bmatrix} -1 & & & 0 \\ & -1 & & 0 \\ & & & 0 \\ 0 & & & c^2 \alpha(x_1, x_3) \end{bmatrix}$$

$J_x = -\sum_\nu m g_{1\nu} \frac{dx^\nu}{ds} = -\sum_\nu m g_{1\nu} \frac{dx^\nu}{dt} \frac{dt}{ds} = \frac{m x_1}{\sqrt{c^2 - q^2}}$

$K_x = -\frac{1}{2} m \frac{\partial g_{44}}{\partial x_1} \frac{dx^4}{ds} \cdot \frac{dx^4}{dt} = -\frac{1}{2} m \frac{\partial c^2}{\partial x_1} \frac{1}{\sqrt{c^2 - q^2}} = -m c \frac{\partial c}{\partial x_1} \frac{1}{\sqrt{c^2 - q^2}}$

all others constant $\frac{1}{ds} = \frac{1}{dt}$

As before

§4 motion of continuously distributed incoherent masses

Pressureless dust



Element of mass m

From section §2, we have expressions for momentum J , Force K , Energy E & Equations of motion



Re-express these equations in terms of densities of momentum, energy, force

$$\sum_{\mu\nu} \frac{\partial}{\partial x^\nu} (\sqrt{g} g_{\mu\alpha} \Theta_{\mu\nu}) - \frac{1}{2} \sum_{\mu\nu} \sqrt{g} \frac{\partial g_{\mu\nu}}{\partial x^\alpha} \Theta_{\mu\nu} = 0$$

accumulation energy-momentum

gravitational force density

where $\Theta_{\mu\nu} = \rho_0 \frac{dx^\mu}{ds} \frac{dx^\nu}{ds}$

contravariant stress-energy tensor
covariant form is $T_{\mu\nu}$
i.e. Einstein "raises" indices
 $T \rightarrow \Theta$

x -momentum density is $-\sqrt{g} g_{\mu\nu} \Theta_{\nu\alpha}$
Energy density is $-\sqrt{g} g_{\mu\nu} \Theta_{\mu\alpha}$

Law of Energy-momentum conservation

• Grossmann. Math Section. §4: Above law corresponds to

covariant divergence $\Theta_{\mu\nu} = 0$... special form possible since $\Theta_{\mu\nu} = \Theta_{\nu\mu}$ (symmetric) } Natural generalization of result in special relativity

• Generalize law:

For all systems, this is the law of energy-momentum conservation.

Reminder: Stress-Energy Tensor in Special Relativity

In any medium

$$c=1$$

Energy conservation

Momentum conservation

energy density \rightarrow $\frac{\partial w}{\partial t}$ energy flux \rightarrow $\nabla \cdot \underline{s}$

$$\frac{\partial w}{\partial t} + \nabla \cdot \underline{s} = 0$$

momentum density \rightarrow $\frac{\partial g}{\partial t}$ stress tensor of medium \rightarrow $\nabla \cdot \underline{t}$

$$\frac{\partial g}{\partial t} + \nabla \cdot \underline{t} = 0$$

$$\begin{aligned} \frac{\partial w}{\partial t} + \frac{\partial s_x}{\partial x} + \frac{\partial s_y}{\partial y} + \frac{\partial s_z}{\partial z} &= 0 \\ \frac{\partial g_x}{\partial t} + \frac{\partial t_{xx}}{\partial x} + \frac{\partial t_{xy}}{\partial y} + \frac{\partial t_{xz}}{\partial z} &= 0 \\ \frac{\partial g_y}{\partial t} + \frac{\partial t_{yx}}{\partial x} + \frac{\partial t_{yy}}{\partial y} + \frac{\partial t_{yz}}{\partial z} &= 0 \\ \frac{\partial g_z}{\partial t} + \frac{\partial t_{zx}}{\partial x} + \frac{\partial t_{zy}}{\partial y} + \frac{\partial t_{zz}}{\partial z} &= 0 \end{aligned}$$

$$\Rightarrow \frac{\partial T_{ik}}{\partial x_k} = 0$$

$x_k = t, x, y, z$ energy flux

T_{ik}	w	s_x	s_y	s_z
	g_x	t_{xx}	t_{xy}	t_{xz}
	g_y	t_{yx}	t_{yy}	t_{yz}
	g_z	t_{zx}	t_{zy}	t_{zz}


momentum density

Symmetry of T_{ik} expresses version of $E=mc^2$:
 momentum density = energy flux

Simplest stress energy tensor for dust
 dust trajectories

$$T_{ik} = \rho v_i v_k$$

In momentarily rest frame of dust, velocity = 0 $\Rightarrow v_i = (1, 0, 0, 0)$




$$T_{ik} = \begin{bmatrix} \rho & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

65 The Differential Equations of the Gravitational Field

The Problem posed:

Poisson's equation
in Newtonian
theory
 $\Delta\phi = 4\pi k\rho$



Equations that determine
how gravitational field $g_{\mu\nu}$
generated by sources &
spread through space

ten components
in $g_{\mu\nu}$
 \therefore seek 10
equations
i.e. second
rank tensor
equations

$$\kappa \Theta_{\mu\nu} = \Gamma_{\mu\nu}$$

stress-
energy tensor
natural
mass source
term

"Gravitation tensor"
constructed from
 $g_{\mu\nu}, \frac{\partial g_{\mu\nu}}{\partial x^\lambda}, \frac{\partial^2 g_{\mu\nu}}{\partial x^\lambda \partial x^\lambda}$

2nd derivatives

Analogous to $\Delta\phi$

Form expected for $\Gamma_{\mu\nu}$:

new " $g^{\mu\nu}$ "

$$\Gamma_{\mu\nu} = \sum_{\alpha\beta} \frac{\partial}{\partial x^\alpha} \left(\gamma_{\alpha\beta} \frac{\partial \delta_{\mu\nu}}{\partial x^\beta} \right) +$$

terms in $\frac{\partial \delta_{\mu\nu}}{\partial x^\alpha}$ that
vanish in weak field

"Core form"

$g_{\mu\nu} \approx \begin{bmatrix} -1 & & \\ & -1 & \\ & & -c^2 \end{bmatrix}$ weak field

In weak field, $g_{\mu\nu}$
differs from $\gamma_{\mu\nu}$ in terms
that are ϵ small

$$- \frac{\partial^2 \delta_{\mu\nu}}{\partial x_1^2} - \frac{\partial^2 \delta_{\mu\nu}}{\partial x_2^2} - \frac{\partial^2 \delta_{\mu\nu}}{\partial x_3^2} + \frac{1}{c^2} \frac{\partial^2 \delta_{\mu\nu}}{\partial x_4^2}$$

so terms like

$$\sum_{\alpha\beta} \frac{\partial g_{\alpha\beta}}{\partial x^\mu} \cdot \frac{\partial \delta_{\alpha\beta}}{\partial x^\nu} \sim \epsilon^2$$

are expected


In static fields only non-zero
term is $\left(\frac{\partial^2}{\partial x_1^2} + \frac{\partial^2}{\partial x_2^2} + \frac{\partial^2}{\partial x_3^2} \right) \delta_{44}$

$$= \Delta(c^2) \approx 2c\Delta c$$

Einstein cannot find a generally tensor for $\Gamma_{\mu\nu}$!

Attempts:

(1) Form derivatives of $g_{\mu\nu}$ using covariant derivative



covariant derivative $g_{\mu\nu} \equiv 0$
condition using it will place no constraint on $g_{\mu\nu}$

(2) Grossmann (mathematical part. §4.2)

Build $\Gamma_{\mu\nu}$ from "Riemann differential tensor"

$$R_{iklm} = (ik, lm) = \frac{1}{2} \left(\frac{\partial^2 g_{im}}{\partial x_k \partial x_l} + \frac{\partial^2 g_{kl}}{\partial x_i \partial x_m} - \frac{\partial^2 g_{il}}{\partial x_k \partial x_m} - \frac{\partial^2 g_{mk}}{\partial x_l \partial x_i} \right) + \sum_{p,q} \chi_{pq} \left(\begin{matrix} im \\ p \end{matrix} \right) \begin{matrix} kl \\ \omega \end{matrix} - \begin{matrix} il \\ p \end{matrix} \begin{matrix} km \\ \omega \end{matrix}$$



contract over 2 indices

$$G_{im} = \sum_k \{ ik, km \}$$

Now called the "Ricci tensor"
 $G_{im} = 0$ are the source free equations of the final theory

"It turns out, however, that in the special case of infinitely weak, static gravitational fields this tensor does not reduce to the expression $\Delta\phi$."



What went wrong?
 Standard view circa 1980 (e.g. Pais biography)

$$G_{im} = \frac{1}{2} \gamma^{kl} \left[\frac{\partial^2 g_{im}}{\partial x_k \partial x_l} + \frac{\partial^2 g_{kl}}{\partial x_i \partial x_m} - \frac{\partial^2 g_{il}}{\partial x_k \partial x_m} - \frac{\partial^2 g_{mk}}{\partial x_l \partial x_i} \right] + \text{Terms quadratic in } \frac{\partial g_{uv}}{\partial x^\alpha}$$

Vanish as ϵ^2 small in weak field

weak field
 \downarrow
 $\sum \left(\frac{\partial^2}{\partial x^i \partial x^i} + \dots \right) g_{im}$
 \downarrow
 $\Delta \phi$

How can we eliminate these terms so G_{im} reduces to $\Delta \phi$?

Expression valid for ALL coordinate systems

Expression valid only in linearly related coordinate systems

so reduction can ONLY succeed if we restrict our coordinate systems.

(New) standard device = specify four conditions to fix coordinate system

e.g.
 "harmonic condition"
 $\gamma_{kl} \left[\begin{smallmatrix} kl \\ i \end{smallmatrix} \right] = 0$ \Rightarrow choice of harmonic coordinates
 3 second derivative terms vanish

plausibility. coordinate system = 4 arbitrary scalar fields.
 \uparrow 4 conditions use up freedom in this arbitrariness

Supposition: Einstein did not recognize this freedom!

Stackel: Perhaps not so simple!
1979

Grossman reports failure of G_{im} for weak STATIC fields:

$$\boxed{G_{im} = 0}$$

→ solve for weak static field of Einstein's 1913 theory

$$g_{uv} = \begin{bmatrix} -1 & 0 & 0 & 0 \\ 0 & -1 & 0 & 0 \\ 0 & 0 & -1 & 0 \\ 0 & 0 & 0 & 1+\phi \end{bmatrix}$$

Natural assumption

- Principle of equivalence
- Solution to $\square g_{uv} = kT_{uv}$ for static dust cloud
- G_{44} only enters in equation of motion of a mass point in weak field.

$$G_{44} = \frac{1}{2} \Delta \phi = 0$$

$$G_{ij} = -\frac{1}{2} \frac{\partial^2 \phi}{\partial x_i \partial x_j} = 0 \quad i, j = 1, 2, 3$$

Disaster!

ϕ varies as linear function of coordinates only.

BUT... in final theory, weak fields are

$$g_{uv} \approx \begin{bmatrix} -1-\phi & 0 & 0 & 0 \\ 0 & -1-\phi & 0 & 0 \\ 0 & 0 & -1-\phi & 0 \\ 0 & 0 & 0 & 1+\phi \end{bmatrix}$$

i.e. weak static fields are not spatially flat.

Einstein's solution : Give up. constraint of energy-momentum conservation already all but fixes field equations
 Leave covariance open!

Lesson of 1912

Einstein illustrates his method = Trivial case of electrostatics
 Field equation $\partial_i \partial_i \phi = \rho$, ($\partial_i = \frac{\partial}{\partial x_i}$)

Force density on charges = $(\partial_i \phi) \rho$ $\xrightarrow{\text{substitute}}$ $\partial_i \phi (\partial_k \partial_k \phi) = \partial_k (\partial_i \phi \partial_k \phi - \frac{1}{2} \delta_{ik} (\partial_m \phi \partial_m \phi))$

Divergence of stress tensor. must get this else action/reaction violated --- lesson of 1912

This much is just an identity. Existence of stress tensor comp. place no further constraint on ϕ after field equation fixed!

DISCOVER field equation by reversing calculation.

① Find identities in ϕ of form.

$$\partial_i \phi \left(\begin{smallmatrix} \text{something} \\ \text{in } \partial_k \partial_k \phi \end{smallmatrix} \right) = \partial_i \left(\begin{smallmatrix} \text{something in } \partial_k \phi \\ \text{that looks like stress tensor} \end{smallmatrix} \right) \text{ e.g.}$$

(2) Read off Field term = ρ in field equation

\downarrow
 $\partial_k \partial_k \phi$ is we use the above identity!

Einstein generates the Entwurf Gravitational Field Equations

seek identity of the form: From force density term in energy-momentum conservation law

$$\frac{\partial}{\partial x^\alpha} \left(\begin{array}{l} \text{something that looks} \\ \text{like a grav. field} \\ \text{stress tensor} \end{array} \right) = \frac{1}{2} \sum_{\mu\nu} \sqrt{g} \frac{\partial g_{\mu\nu}}{\partial x^\alpha} \left(\begin{array}{l} \text{something in 1st, 2nd} \\ \text{derivatives of } g_{\mu\nu} \\ \text{of core form} \end{array} \right)$$

Grossmann (mathematical part 54) supplies identity:

$$\sum_{\alpha\beta\gamma\rho} \frac{\partial}{\partial x^\alpha} \left(\sqrt{-g} \delta_{\alpha\beta} \frac{\partial \delta_{\gamma\rho}}{\partial x^\beta} \cdot \frac{\partial g_{\gamma\rho}}{\partial x^\alpha} \right) - \frac{1}{2} \sum_{\alpha\beta\gamma\rho} \frac{\partial}{\partial x^\alpha} \left(\sqrt{-g} \delta_{\alpha\beta} \frac{\partial \delta_{\gamma\rho}}{\partial x^\alpha} \cdot \frac{\partial g_{\gamma\rho}}{\partial x^\beta} \right) \left. \vphantom{\sum} \right\} \text{Read off stress-tens-ene}$$

$$= \sum_{\mu\nu} \sqrt{g} \frac{\partial g_{\mu\nu}}{\partial x^\alpha} \left\{ \sum_{\alpha\beta} \frac{1}{\sqrt{g}} \frac{\partial}{\partial x^\alpha} \left(\delta_{\alpha\beta} \sqrt{g} \frac{\partial \delta_{\mu\nu}}{\partial x^\beta} \right) - \sum_{\alpha\beta\gamma\rho} \delta_{\alpha\beta} g_{\gamma\rho} \frac{\partial \delta_{\mu\epsilon}}{\partial x^\alpha} \frac{\partial \delta_{\nu\rho}}{\partial x^\beta} \right. \\ \left. + \frac{1}{2} \sum_{\alpha\beta\gamma\rho} \delta_{\alpha\mu} \delta_{\beta\nu} \frac{\partial g_{\gamma\rho}}{\partial x^\alpha} \frac{\partial \delta_{\gamma\rho}}{\partial x^\beta} - \frac{1}{4} \sum_{\alpha\beta\gamma\rho} \delta_{\mu\nu} \delta_{\alpha\beta} \frac{\partial g_{\gamma\rho}}{\partial x^\alpha} \frac{\partial \delta_{\gamma\rho}}{\partial x^\beta} \right\}$$

This is $2 \Gamma_{\mu\nu}$