

## Lecture 22/Chapter 19

### Part 4. Statistical Inference Ch. 19

#### Diversity of Sample Proportions

- Probability versus Inference
- Behavior of Sample Proportions: Example
- Behavior of Sample Proportions: Conditions
- Behavior of Sample Proportions: Rules

## Course Divided into Four Parts (Review)

1. Finding Data in Life: scrutinizing origin of data
2. Finding Life in Data: summarizing data yourself or assessing another's summary
3. Understanding Uncertainty in Life: probability theory (completed)
4. Making Judgments from Surveys and Experiments: statistical inference

## Approach to Inference

- **Step 1 (Chapter 19):** Work *forward*---if we happen to know the population proportion falling in a given category, what behavior can we expect from sample proportions for repeated samples of a given size?
  - **Step 2 (Chapter 20):** Work *backward*---if sample proportion for a sample of a certain size is observed to take a specified value, what can we conclude about the value of the unknown population proportion?
- After covering Steps 1&2 for **proportions**, we'll cover them for **means**.

## Understanding Sample Proportion

- 3 Approaches:
1. Intuition
  2. Hands-on Experimentation
  3. Theoretical Results

We'll find that our **intuition** is consistent with **experimental** results, and both are confirmed by mathematical **theory**.


**Example:** *Intuit Behavior of Sample Proportion*

- **Background:** Population proportion of blue M&M's is  $1/6=0.17$ .
- **Question:** How does sample proportion behave for repeated random samples of size 25 (a teaspoon)?
- **Response:** Summarize by telling
- **Experiment:** sample teaspoons of M&Ms, record sample proportion of blues on sheet **and** in notes (need a calculator)

**Example:** *Intuit Behavior of Sample Proportion*

- **Background:** Population proportion of blue M&M's is  $1/6=0.17$ .

**Note:** The shape of the underlying distribution (sample size 1) will play a role in the shape of sample proportions for various sample sizes.



**Example:** *Intuit Behavior of Sample Proportion*

- **Response:** (continued)
  - **Center:** some sample proportions will be less than 0.17 and others more; the mean of all sample proportions should be \_\_\_\_\_
  - **Spread:** depends on sample size; if we'd sampled only 5, we'd easily get sample proportions ranging from 0 to 0.6 or 0.8. For samples of 25, proportions \_\_\_\_\_
  - **Shape:** proportions close to \_\_\_\_\_ would be most common, and those far from \_\_\_\_\_ increasingly less likely---shape\_\_\_\_\_

**Example:** *Intuit Behavior of Sample Proportion*

- **Background:** Population proportion of blue M&M's is  $1/6=0.17$ .
- **Question:** How does sample proportion behave for repeated random samples of size 75 (a Teaspoon)?
- **Response:** Again, we summarize by telling
  - Now sample Tablespoons of M&Ms, record sample proportion of blues on sheet **and** in notes (need a calculator)

### Example: Intuit Behavior of Sample Proportion

- **Response:** (samples of size 75)
- **Center:** The mean of all sample proportions should be \_\_\_\_\_, regardless of sample size.
- **Spread:** should be \_\_\_\_\_ than what it would be for samples of size 25.
- **Shape:** should bulge more close to 0.17, taper more at the ends, less right-skewness: it should be \_\_\_\_\_.

### Conditions for Rule of Sample Proportions

- **Randomness** [affects center]
  - Can't be biased for or against certain values
- **Independence** [affects spread]
  - If sampling without replacement, sample should be less than  $1/10$  population size
- **Large enough sample size** [affects shape]
  - Should sample enough to expect at least 5 each in and out of the category of interest.

### Example: Checking Conditions for Rule

- **Background:** Population proportion of blue M&M's is  $1/6=0.17$ . Students repeatedly take random samples of size 1 teaspoon (about 25) and record the proportion that are blue.
- **Question:** Are the 3 Conditions met?
- **Response:**
  1. \_\_\_\_\_
  2. \_\_\_\_\_
  3. \_\_\_\_\_

### Example: Checking Conditions (larger sample)

- **Background:** Population proportion of blue M&M's is  $1/6=0.17$ . Students repeatedly take random samples of size 1 Tablespoon (about 75) and record the proportion that are blue.
- **Question:** Are the 3 Conditions met?
- **Response:**
  1. \_\_\_\_\_
  2. \_\_\_\_\_
  3. \_\_\_\_\_

## Rule for Sample Proportions

- **Center:** The mean of sample proportions equals the true population proportion.
- **Spread:** The standard deviation of sample proportions is standard error =  $\sqrt{\frac{\text{population proportion} \times (1 - \text{population proportion})}{\text{sample size}}}$
- **Shape:** (Central Limit Theorem) The frequency curve of proportions from the various samples is approximately normal.

## Example: Applying Rules for Sample Proportions

- **Background:** Proportion of blue M&Ms is  $1/6=0.17$ .
  - **Question:** What does the Rule tell us about sample proportions that are blue in teaspoons (about 25)?
  - **Response:**
    - Center: the mean of sample proportions will be \_\_\_\_\_
    - Spread: the standard deviation of sample proportions will be \_\_\_\_\_
- standard error = \_\_\_\_\_
- Shape: \_\_\_\_\_

## Example: Applying Rules for Sample Proportions

- **Background:** Proportion of blue M&Ms is  $1/6=0.17$ .
  - **Question:** What does the Rule tell us about sample proportions that are blue in Teaspoons (about 75)?
  - **Response:**
    - Center: the mean of sample proportions will be \_\_\_\_\_
    - Spread: the standard deviation of sample proportions will be \_\_\_\_\_
- standard error = \_\_\_\_\_
- Shape: \_\_\_\_\_

## Empirical Rule (Review)

- For any normal curve, approximately
- 68% of values are within 1 sd of mean
  - 95% of values are within 2 sds of mean
  - 99.7% of values are within 3 sds of mean

### Example: Applying Empirical Rule to M&Ms

□ **Background:** Population proportion of blue M&M's is  $1/6=0.17$ . Students repeatedly take random samples of size 1 Tablespoon (about 75) and record the proportion that are blue.

□ **Question:** What does the Empirical Rule tell us?

□ **Response:**

- 68% of the sample proportions should be within \_\_\_\_\_: in [0.127, 0.213]
- 95% of the sample proportions should be within \_\_\_\_\_: in [0.084, 0.256]
- 99.7% of the sample proportions should be within \_\_\_\_\_: in [0.041, 0.299]

How well did our sampled proportions conform?

### Proportions then Means, Probability then Inference

Next time we'll establish a parallel theory for means, when the variable of interest is quantitative (number on dice instead of color on M&M). After that, we'll

- Perform inference with **confidence intervals**
  - For proportions (Chapter 20)
  - For means (Chapter 21)
- Perform inference with **hypothesis testing**
  - For proportions (Chapters 22&23)
  - For means (Chapters 22&23)