

# CS 441: Propositional Logic

---

PhD. Nils Murrugarra-Llerena  
[nem177@pitt.edu](mailto:nem177@pitt.edu)



## Today's Topic: Propositional Logic

- What is a proposition?
- Logical connectives and truth tables
- Translating between English and propositional logic



## Logic is the basis of all mathematical and analytical reasoning

Given a collection of known truths, logic allows us to deduce new truths

### *Example*

#### Base facts:

If it is raining, I will not go outside

If I am inside, Lisa will stay home

Lisa and I always play video games if we are together during the weekend

Today is a rainy Saturday

**Conclusion:** Lisa and I will play video games today

Logic allows us to advance mathematics through an iterative process of **conjecture** and **proof**

# Propositional logic is a very simple logic

**Definition:** A **proposition** is a precise statement that is either **true** or **false**, but not both.

Examples:

- $2 + 2 = 4$  (**true**)
- All dogs have 3 legs (**false**)
- $x^2 < 0$  (**false**)
- Washington, D.C. is the capital of the USA (**true**)

## Not all statements are propositions

- Eliana is cool
  - “Cool” is a subjective term.
- $x^3 < 0$ 
  - **True** if  $x < 0$ , **false** otherwise.
- Springfield is the capital
  - **True** in Illinois, **false** in Massachusetts.

## We can use logical connectives to build complex propositions

We will discuss the following logical connectives:

- $\neg$  (not)
- $\wedge$  (conjunction / and)
- $\vee$  (disjunction / or)
- $\oplus$  (exclusive disjunction / xor)
- $\rightarrow$  (implication)
- $\leftrightarrow$  (biconditional)

# Negation

The **negation** of a proposition is **true** iff the proposition is **false**

What we know

What we want to know

One row for each possible value of "what we know"

$p$	$\neg p$

The truth table for negation

*(I'll sometimes use  $\top$  and  $\perp$ )*

## Negation Examples

Negate the following propositions

- Today is Monday
- $21 * 2 = 42$


What is the truth value of the following propositions

- $\neg$ (9 is a prime number)
- $\neg$ (Pittsburgh is in Pennsylvania)



# Conjunction

The **conjunction** of two propositions is true iff both propositions are true



p	q	$p \wedge q$
T	T	
T	F	
F	T	
F	F	

The truth table for conjunction

***$2^2 = 4$  rows since we know both  $p$  and  $q$ !***

## Disjunction

The **disjunction** of two propositions is true iff *at least one* proposition is true

p	q	$p \vee q$
T	T	
T	F	
F	T	
F	F	

The truth table for disjunction

## Conjunction and disjunction examples

*This symbol means “is defined as”  
or “is equivalent to”  
(sometimes seen as  $\triangleq$ )*

Let:

- $p \equiv x^2 \geq 0$  True
- $q \equiv$  A lion weighs less than a mouse False
- $r \equiv 10 < 7$  False
- $s \equiv$  Pittsburgh is located in Pennsylvania True

What are the truth values of these expressions:

- $p \wedge q$
- $p \wedge s$
- $p \vee q$
- $q \vee r$

## In-class Exercises

**Problem 1:** Let  $p \equiv 2+2=5$ ,  $q \equiv$  eagles can fly,  $r \equiv 1=1$ . Determine the value for each of the following:

- $p \wedge q$
- $\neg p \vee q$
- $p \vee (q \wedge r)$
- $(p \vee q) \wedge (\neg r \vee \neg p)$

## Exclusive or (XOR)

The **exclusive or** of two propositions is true iff *exactly one* proposition is true

p	q	$p \oplus q$
T	T	
T	F	
F	T	
F	F	

The truth table for **exclusive or**

**Note:** Exclusive or is typically used to natural language to identify *choices*. For example, “You may have a soup or salad with your entree.”

# Implication

The **implication**  $p \rightarrow q$  is **false** if  $p$  is **true**, and  $q$  is **false**;  $p \rightarrow q$  is **true** otherwise

## Terminology

- $p$  is called the hypothesis
- $q$  is called the conclusion

$p$	$q$	$p \rightarrow q$
T	T	
T	F	
F	T	
F	F	

The truth table for **implication**

## Implication (cont.)

The implication  $p \rightarrow q$  can be read in a number of (equivalent) ways:

- If  $p$  then  $q$
- $p$  only if  $q$
- $p$  is sufficient for  $q$
- $q$  whenever  $p$

## Implication examples

Let:

- $p \equiv$  Jane gets a 100% on her final exam
- $q \equiv$  Jane gets an A on her final exam

What are the truth values of these implications:

- $p \rightarrow q$  -----
- $q \rightarrow p$



## Other conditional statements

Given an implication  $p \rightarrow q$ :

- $q \rightarrow p$  is its **converse**
- $\neg q \rightarrow \neg p$  is its **contrapositive**
- $\neg p \rightarrow \neg q$  is its **inverse**



Why might this be useful?

**Note:** An **implication** and its **contrapositive** *always* have the same truth value

## Biconditional

The biconditional  $p \leftrightarrow q$  is true if and only if  $p$  and  $q$  assume the same truth value

$p$	$q$	$p \leftrightarrow q$
T	T	
T	F	
F	T	—
F	F	

The truth table for the biconditional

**Note:** The biconditional statement  $p \leftrightarrow q$  is often read as “ $p$  if and only if  $q$ ” or “ $p$  is a necessary and sufficient condition for  $q$ .”



## Like mathematical operators, logical operators are assigned precedence levels

1. Negation
  - $\neg q \vee r$  means  $(\neg q) \vee r$ , not  $\neg(q \vee r)$
2. Conjunction
3. Disjunction
  - $q \wedge r \vee s$  means  $(q \wedge r) \vee s$ , not  $q \wedge (r \vee s)$
4. Implication
  - $q \wedge r \rightarrow s$  means  $(q \wedge r) \rightarrow s$ , not  $q \wedge (r \rightarrow s)$
5. Biconditional

In general, we will use **parentheses** to disambiguate and to override precedence rules.

## In-class Exercises

**Problem 2:** Show that an implication  $p \rightarrow q$  and its contrapositive  $\neg q \rightarrow \neg p$  always have the same value

- **Hint:** Construct two truth tables

**Problem 3:** Construct the truth table for the compound proposition  $p \wedge (\neg q \vee r) \rightarrow s$

## English sentences can often be translated into propositional sentences


But *why* would we do that?



Philosophy and epistemology



Reasoning about law



Verifying complex system specifications

## Example #1

**Example:** You can see an R-rated movie **only if** you are over 17 **or** you are accompanied by your legal guardian.

Let:

Find logical connectives



Translate fragments



Create logical expression



## Example #2

**Example:** You can have free coffee **if** you are a senior citizen **and** it is a Tuesday

**Let:**



## Example #3

**Example:** If you are under 17 and are not accompanied by your legal guardian, then you cannot see the R-rated movie.

**Let:**

**Note:** The above translation is the contrapositive of the translation from example 1!

## Logic also helps us understand bitwise operations

- Computers represent data as sequences of bits
  - e.g., 0101 1101 1010 1111
- Bitwise logical operations are often used to manipulate these data
- If we treat 1 as **true** and 0 as **false**, our logic truth tables tell us how to carry out bitwise logical operations

## Bitwise logic examples

$$\begin{array}{r} \wedge \\ \hline 1010\ 1110 \\ 1110\ 1010 \\ \hline \end{array}$$

$$\begin{array}{r} \vee \\ \hline 1010\ 1110 \\ 1110\ 1010 \\ \hline \end{array}$$

$$\begin{array}{r} \oplus \\ \hline 1010\ 1110 \\ 1110\ 1010 \\ \hline \end{array}$$

## In-class Exercises

**Problem 4:** Translate the following sentences

- On Top Hat

**Problem 5:** Solve the following bitwise problems

$$\begin{array}{r} \oplus \\ \hline 1011\ 1000 \\ 1010\ 0110 \end{array}$$

$$\begin{array}{r} \wedge \\ \hline 1011\ 1000 \\ 1010\ 0110 \end{array}$$

## Final Thoughts

- Propositional logic is a simple logic that allows us to reason about a variety of concepts
- In recitation:
  - More examples and practice problems
  - Be sure to attend!
- Next:
  - Logic puzzles and propositional equivalence
  - Please read sections 1.2 and 1.3
    - In general: do the assigned reading!