

**Exercise (2.1.3).** Is the sequence  $\left\{\frac{(-1)^n}{2^n}\right\}$  convergent? If so, what is the limit?

Answer: The sequence converges to 0.

*Proof.* Note that, for all  $n \in \mathbb{N}$ ,

$$-\frac{1}{n} \leq \frac{(-1)^n}{2^n} \leq \frac{1}{n}.$$

Since  $\left\{\pm\frac{1}{n}\right\}$  converges to 0,  $\left\{\frac{(-1)^n}{2^n}\right\}$  converges to 0 by the Squeeze Theorem.  $\square$

**Exercise (2.1.15).** Let  $\{x_n\}$  be a sequence defined by

$$x_n := \begin{cases} n & \text{if } n \text{ is odd,} \\ 1/n & \text{if } n \text{ is even.} \end{cases}$$

a) Is the sequence bounded? (prove or disprove)

b) Is there a convergent subsequence? If so, find it.

*Solution.* a)  $\{x_n\}$  is not bounded.

*Proof.* For any  $M \in \mathbb{R}$ , by the Archimedean property,  $\exists n \in \mathbb{N}$  such that  $n > M$ . If  $n$  is odd,  $x_n = n > M$ . If  $n$  is even,  $n + 1$  is odd and  $x_{n+1} = n + 1 > M$ . Hence,  $\{x_n\}$  cannot be bounded.  $\square$

b) Let  $n_i = 2i$  for all  $i \in \mathbb{N}$ . Then  $\{x_{n_i}\}_i = \{1/2i\}_i$ , which converges to zero.  $\square$

**Exercise (2.2.7).** True or false, prove or find a counterexample. If  $(x_n)$  is a sequence such that  $(x_n^2)$  converges, then  $(x_n)$  converges.

*Solution.* This is false. One counterexample is given by  $x_n := (-1)^n$ .  $\square$

**Exercise (2.3.7).** Let  $\{x_n\}$  and  $\{y_n\}$  be bounded sequences.

a) Show that  $\{x_n + y_n\}$  is bounded.

b) Show that

$$(\liminf_{n \rightarrow \infty} x_n) + (\liminf_{n \rightarrow \infty} y_n) \leq \liminf_{n \rightarrow \infty} (x_n + y_n).$$

c) Find explicit  $\{x_n\}$  and  $\{y_n\}$  such that

$$(\liminf_{n \rightarrow \infty} x_n) + (\liminf_{n \rightarrow \infty} y_n) < \liminf_{n \rightarrow \infty} (x_n + y_n).$$

*Proof.* a) Since  $\{x_n\}$  and  $\{y_n\}$  are bounded, there are numbers  $B_1$  and  $B_2$  such that for all  $n \in \mathbb{N}$ ,

$$|x_n| \leq B_1 \quad \text{and} \quad |y_n| \leq B_2.$$

Then, for all  $n \in \mathbb{N}$ ,

$$\begin{aligned} |x_n + y_n| &\leq |x_n| + |y_n| \\ &\leq B_1 + B_2. \end{aligned}$$

Hence,  $\{x_n + y_n\}$  is a bounded sequence.

b) Note that, for any  $n \in \mathbb{N}$  and  $j \geq n$ ,

$$x_j + y_j \geq \inf\{x_k : k \geq n\} + \inf\{y_k : k \geq n\}.$$

Hence,  $\inf\{x_k : k \geq n\} + \inf\{y_k : k \geq n\}$  is a lower bound of the set  $\{x_k + y_k : k \geq n\}$ . Therefore, for all  $n \in \mathbb{N}$ ,

$$\inf\{x_k : k \geq n\} + \inf\{y_k : k \geq n\} \leq \inf\{x_k + y_k : k \geq n\}.$$

Taking the limit as  $n \rightarrow \infty$  of both sides of this inequality, we obtain

$$(\liminf_{n \rightarrow \infty} x_n) + (\liminf_{n \rightarrow \infty} y_n) \leq \liminf_{n \rightarrow \infty} (x_n + y_n).$$

c) Let  $\{x_n\} = \{(-1)^{n+1}\}$  and  $\{y_n\} = \{(-1)^n\}$ . Then  $\{x_n + y_n\} = \{0\}$ , for all  $n \in \mathbb{N}$ . Hence,

$$\begin{aligned} (\liminf_{n \rightarrow \infty} x_n) + (\liminf_{n \rightarrow \infty} y_n) &= (-1) + (-1) \\ &= -2 \\ &< 0 \\ &= \liminf_{n \rightarrow \infty} (x_n + y_n). \end{aligned}$$

□

**Exercise.** Show that if  $(x_n)_{n \in \mathbb{N}}$  is a convergent sequence then every subsequence of  $(x_n)_{n \in \mathbb{N}}$  is also convergent. Moreover if

$$x := \lim_{n \rightarrow \infty} x_n$$

then for any subsequence  $(x_{n_i})_i$ ,

$$x = \lim_{i \rightarrow \infty} x_{n_i}$$

*Proof.* Assume that  $(x_n)$  converges to  $x$  as  $n \rightarrow \infty$ . Let  $(x_{n_i})$  be any subsequence.

Since  $(x_n)$  converges, given  $\epsilon > 0$ ,  $\exists N \in \mathbb{N}$  such that  $n \geq N \Rightarrow |x_n - x| < \epsilon$ . Then, for any  $i \in \mathbb{N}$  such that  $i \geq N$ , since  $n_i \geq i \geq N$ ,  $|x_{n_i} - x| < \epsilon$ . Hence,  $(x_{n_i})$  also converges to  $x$ .  $\square$

**Exercise.** Let  $(x_n)_{n \in \mathbb{N}}$  be a sequence and assume one of the following properties:

a) there is some  $x$  such that any subsequence  $(x_{n_i})_i$  contains another subsequence  $(x_{n_{i_j}})_{j \in \mathbb{N}}$  which is convergent to  $x$ .

b) any subsequence  $(x_{n_i})_i$  contains another subsequence  $(x_{n_{i_j}})_{j \in \mathbb{N}}$  which is convergent (a priori not necessarily to the same  $x$ )

Show in which cases  $(x_n)_n$  is convergent. Give a counterexample for the other cases.

*Solution.* Assume that there is some  $x$  such that any subsequence  $(x_{n_i})_i$  contains another subsequence  $(x_{n_{i_j}})_{j \in \mathbb{N}}$  which converges to  $x$ .

If  $(x_n)$  is not convergent, there must be some  $\epsilon > 0$  and a subsequence  $(x_{n_i})$  such that

$$|x_{n_i} - x| > \epsilon \quad \forall i \in \mathbb{N}.$$

But then this  $(x_{n_i})_i$  cannot have another subsequence  $(x_{n_{i_j}})_{j \in \mathbb{N}}$  that converges to  $x$ , because still

$$|x_{n_{i_j}} - x| > \epsilon \quad \forall i \in \mathbb{N}.$$

This is a contradiction. Hence,  $\lim_{n \rightarrow \infty} x_n = x$ .

b) Assume that any subsequence  $(x_{n_i})_i$  contains another subsequence  $(x_{n_{i_j}})_{j \in \mathbb{N}}$  which is convergent (a priori not necessarily to the same  $x$ ). In this case,  $(x_n)$  does not have to converge.

Counterexample:  $x_n := (-1)^n$ . Not convergent, but any subsequence has another subsequence which is convergent.

$\square$

**Exercise** (bonus). *Find the following limit. Show all work.*

$$\lim_{n \rightarrow \infty} \left( \frac{1}{\sqrt{n^2+1}} + \frac{1}{\sqrt{n^2+2}} + \dots + \frac{1}{\sqrt{n^2+2n}} \right)$$

*Proof.* Set

$$x_n := \left( \frac{1}{\sqrt{n^2+1}} + \frac{1}{\sqrt{n^2+2}} + \dots + \frac{1}{\sqrt{n^2+2n}} \right)$$

Then

$$\frac{2n}{\sqrt{n^2+2n}} \geq x_n \leq \frac{2n}{\sqrt{n^2+1}}$$

Now

$$\lim_{n \rightarrow \infty} \frac{2n}{\sqrt{n^2+2n}} = 2 \lim_{n \rightarrow \infty} \frac{n}{n} \frac{1}{\sqrt{1+\frac{2}{n}}} = 2.$$

and

$$\lim_{n \rightarrow \infty} \frac{2n}{\sqrt{n^2+1}} = 2 \lim_{n \rightarrow \infty} \frac{n}{n} \frac{1}{\sqrt{1+\frac{1}{n^2}}} = 2.$$

By the squeeze theorem

$$\lim_{n \rightarrow \infty} x_n = 2.$$

□